



Journal of Frontiers in Multidisciplinary Research

The Effects of Concussion Management Protocols in Contact Sports

Imelda T Angeles

College of Commerce and Business Administration University of Santo Tomas, Manila, Philippines

* Corresponding Author: **Imelda T Angeles**

Article Info

E-ISSN: 3050-9726

P-ISSN: 3050-9718

Volume: 04

Issue: 01

Jan-Jun 2023

Received: 05-02-2023;

Accepted: 04-03-2023

Published: 20-04-2023

Page No: 11-13

Abstract

Concussions are a significant concern in contact sports due to their potential to cause long-term neurological damage. Effective concussion management protocols are essential to mitigate these risks and ensure the safety of athletes. This article explores the effects of concussion management protocols in contact sports, examining their development, implementation, and outcomes. The study reviews existing literature, analyzes case studies, and evaluates the effectiveness of various protocols. The findings suggest that well-designed and consistently applied concussion management protocols can significantly reduce the incidence and severity of concussions, improve recovery times, and enhance overall athlete safety. However, challenges such as compliance, education, and resource allocation remain. This article concludes with recommendations for improving concussion management in contact sports.

Keywords: Concussion, Contact Sports, Management Protocols, Athlete Safety, Neurological Damage, Recovery

Introduction

Concussions are a type of traumatic brain injury (TBI) that occur when a blow to the head or body causes the brain to move rapidly within the skull. This movement can lead to chemical changes in the brain and, in some cases, damage to brain cells. Concussions are particularly prevalent in contact sports such as football, rugby, hockey, and soccer, where collisions and impacts are common. The short-term effects of concussions can include headaches, dizziness, confusion, and memory problems, while long-term effects may involve chronic traumatic encephalopathy (CTE), depression, and cognitive decline.

Given the potential severity of these outcomes, the development and implementation of effective concussion management protocols are critical. These protocols are designed to identify, assess, and manage concussions in a way that minimizes risks and promotes safe recovery. This article aims to provide a comprehensive overview of the effects of concussion management protocols in contact sports, covering their development, implementation, and outcomes.

Materials and Methods

This study employs a mixed-methods approach, combining a review of existing literature with an analysis of case studies and empirical data. The literature review focuses on peer-reviewed articles, clinical guidelines, and policy documents related to concussion management in contact sports. The case studies include examples from professional, collegiate, and youth sports organizations that have implemented concussion management protocols. Empirical data were collected from surveys and interviews with athletes, coaches, and medical professionals.

The study evaluates the effectiveness of various concussion management protocols, including the use of standardized assessment tools, return-to-play guidelines, and educational programs. The analysis considers factors such as compliance rates, recovery times, and long-term outcomes for athletes. The study also examines the challenges and barriers to effective concussion management, including issues related to education, resources, and cultural attitudes toward injury in sports.

Results

The findings of this study indicate that concussion management protocols have a significant impact on athlete safety and

recovery. The use of standardized assessment tools, such as the Sport Concussion Assessment Tool (SCAT), has improved the accuracy of concussion diagnosis and the consistency of care. Return-to-play guidelines, which typically involve a graduated progression of physical activity, have been shown to reduce the risk of re-injury and promote safer recovery. Educational programs aimed at athletes, coaches, and parents have also been effective in increasing awareness of concussion risks and the importance of proper management. Case studies from professional sports leagues, such as the NFL and NHL, demonstrate that organizations with comprehensive concussion management protocols have lower rates of concussion-related complications and better long-term outcomes for athletes.

However, the study also identifies several challenges to effective concussion management. Compliance with protocols can be inconsistent, particularly at the youth and amateur levels where resources and oversight may be limited. Cultural attitudes toward injury in sports, such as the "tough it out" mentality, can also hinder the proper implementation of concussion management protocols. Additionally, there is a need for ongoing education and training to ensure that all stakeholders are aware of the latest guidelines and best practices.

Discussion

The results of this study highlight the importance of well-designed and consistently applied concussion management protocols in contact sports. The use of standardized assessment tools and return-to-play guidelines has been shown to improve outcomes for athletes, reducing the risk of re-injury and promoting safer recovery. Educational programs have also played a crucial role in increasing awareness and understanding of concussion risks.

However, the study also underscores the need for continued efforts to address the challenges and barriers to effective concussion management. Compliance with protocols remains a significant issue, particularly at the youth and amateur levels. Efforts to improve compliance should focus on increasing resources and oversight, as well as changing cultural attitudes toward injury in sports. Ongoing education and training are also essential to ensure that all stakeholders are equipped with the knowledge and skills needed to implement concussion management protocols effectively.

The study also highlights the importance of interdisciplinary collaboration in concussion management. Medical professionals, coaches, athletes, and parents all play a role in ensuring the safety and well-being of athletes. Effective communication and coordination among these stakeholders are critical to the success of concussion management protocols.

Conclusion

Concussion management protocols are a vital component of athlete safety in contact sports. The findings of this study demonstrate that well-designed and consistently applied protocols can significantly reduce the incidence and severity of concussions, improve recovery times, and enhance overall athlete safety. However, challenges such as compliance, education, and resource allocation remain.

To address these challenges, it is essential to continue developing and refining concussion management protocols, with a focus on increasing compliance, improving education and training, and fostering interdisciplinary collaboration. By

doing so, we can ensure that athletes at all levels of contact sports are protected from the potentially devastating effects of concussions.

References

1. McCrory P, Meeuwisse W, Dvořák J, et al. Consensus statement on concussion in sport—the 5th international conference on concussion in sport held in Berlin, October 2016. *British Journal of Sports Medicine*. 2017;51(11):838-47.
2. Giza CC, Hovda DA. The new neurometabolic cascade of concussion. *Neurosurgery*. 2014;75(Suppl 4):S24-S33.
3. Harmon KG, Drezner JA, Gammons M, et al. American Medical Society for Sports Medicine position statement: Concussion in sport. *Clinical Journal of Sport Medicine*. 2013;23(1):1-18.
4. Broglio SP, Cantu RC, Gioia GA, et al. National Athletic Trainers' Association position statement: Management of sport concussion. *Journal of Athletic Training*. 2014;49(2):245-65.
5. Kerr ZY, Register-Mihalik JK, Marshall SW, et al. Disclosure and non-disclosure of concussion and concussion symptoms in athletes: Review and application of the socio-ecological framework. *Brain Injury*. 2014;28(8):1009-21.
6. Chrisman SP, Rivara FP, Schiff MA, et al. Risk factors for concussive symptoms 1 week or longer in high school athletes. *Brain Injury*. 2013;27(1):1-9.
7. Guskiewicz KM, Weaver NL, Padua DA, et al. Epidemiology of concussion in collegiate and high school football players. *American Journal of Sports Medicine*. 2000;28(5):643-50.
8. Meehan WP, d'Hemecourt P, Comstock RD. High school concussions in the 2008-2009 academic year: Mechanism, symptoms, and management. *American Journal of Sports Medicine*. 2010;38(12):2405-9.
9. Register-Mihalik JK, Guskiewicz KM, McLeod TC, et al. Knowledge, attitude, and concussion-reporting behaviors among high school athletes: A preliminary study. *Journal of Athletic Training*. 2013;48(5):645-53.
10. Kerr ZY, Marshall SW, Harding HP, et al. Nine-year risk of depression diagnosis increases with increasing self-reported concussions in retired professional football players. *American Journal of Sports Medicine*. 2012;40(10):2206-12.
11. Guskiewicz KM, Marshall SW, Bailes J, et al. Association between recurrent concussion and late-life cognitive impairment in retired professional football players. *Neurosurgery*. 2005;57(4):719-26.
12. McKee AC, Cantu RC, Nowinski CJ, et al. Chronic traumatic encephalopathy in athletes: Progressive tauopathy after repetitive head injury. *Journal of Neuropathology & Experimental Neurology*. 2009;68(7):709-35.
13. Stern RA, Riley DO, Daneshvar DH, et al. Long-term consequences of repetitive brain trauma: Chronic traumatic encephalopathy. *PM&R*. 2011;3(10 Suppl 2):S460-S67.
14. Gavett BE, Stern RA, McKee AC. Chronic traumatic encephalopathy: A potential late effect of sport-related concussive and subconcussive head trauma. *Clinics in Sports Medicine*. 2011;30(1):179-88.
15. Daneshvar DH, Nowinski CJ, McKee AC, et al. The

- epidemiology of sport-related concussion. *Clinics in Sports Medicine*. 2011;30(1):1-17.
16. Collins MW, Kontos AP, Okonkwo DO, et al. Statements of agreement from the targeted evaluation and active management (TEAM) approaches to treating concussion meeting held in Pittsburgh, October 15-16, 2015. *Neurosurgery*. 2016;79(6):912-29.
 17. Elbin RJ, Sufrinko A, Schatz P, et al. Removal from play after concussion and recovery time. *Pediatrics*. 2016;138(3):e20160910.
 18. Kroshus E, Baugh CM, Daneshvar DH, et al. Pressure on sports medicine clinicians to prematurely return collegiate athletes to play after concussion. *Journal of Athletic Training*. 2015;50(9):944-51.
 19. Kerr ZY, Yeargin S, Valovich McLeod TC, et al. Comprehensive coach education and practice contact restriction guidelines result in lower injury rates in youth American football. *Orthopaedic Journal of Sports Medicine*. 2015;3(7):2325967115594578.
 20. Chrisman SP, Quitiquit C, Rivara FP. Qualitative study of barriers to concussive symptom reporting in high school athletics. *Journal of Adolescent Health*. 2013;52(3):330-5.
 21. Register-Mihalik JK, Linnan LA, Marshall SW, et al. Using theory to understand high school aged athletes' intentions to report sport-related concussion: Implications for concussion education initiatives. *Brain Injury*. 2013;27(7-8):878-86.
 22. Kroshus E, Garnett B, Hawrilenko M, et al. Concussion under-reporting and pressure from coaches, teammates, fans, and parents. *Social Science & Medicine*. 2015;134:66-75.
 23. Kerr ZY, Register-Mihalik JK, Kroshus E, et al. Motivations associated with nondisclosure of self-reported concussions in former collegiate athletes. *American Journal of Sports Medicine*. 2016;44(1):220-5.
 24. Kroshus E, Baugh CM, Stein CJ, et al. Concussion reporting, sex, and conformity to traditional gender norms in young adults. *Journal of Adolescence*. 2017;54:110-9.